

A REVIEW ON *MYRISTICA FRAGRANS* (NUTMEG)

Dr. Anil Kumar*¹, Krishanu Samanta²

*^{1,2}Pharmacy College, Itaura, Chandeshwar, Azamgarh U.P., India.

ABSTRACT

Herbal medicine is the oldest form of healthcare known to mankind. Herbs had been used by all cultures throughout history. *Myristica fragrans* commonly known as 'Nutmeg', is an evergreen tree indigenous to India, Indonesia and Sri Lanka. It is the main source of spices, having a pleasant aromatic fragrance and widely used as flavouring agent and ability to enhance the taste of food. It has been known to provide many medicinal properties due to its complex molecular structure. Many phytoconstituents of *M. fragrans* is responsible for its effect in curing various diseases like antioxidant, antidiarrheal, hypolipidemic, antidepressant, aphrodisiac, anticonvulsant, antimicrobial, memory enhancer and hepatoprotective properties. This review attempts to highlight the available literature traditional uses, and its pharmacological activities of *Myristica fragrans*. This will be helpful to create interest towards Nutmeg and may be useful in developing new formulations with economical value.

KEYWORDS: *Myristica fragrans*, nutmeg, Traditional use. Pharmacological activity.

1. INTRODUCTION

Nature is served as primary source of numerous medicines for thousands of years Medicinal plants/herbs hold a great significant value. Medicinal plants/herbs possess phytochemical constituents that are medically important and are used for developing new drugs. *Myristica fragrance* is most important herbal medicinal plant commonly known as Nutmeg or Jaiphal, belonging to family Myristicaceae. The use of plants for health benefits and other purposes is a very common tradition, which has been in practice since ancient times. Medicinal products derived from the plants are often named as herbal medicines, herbal drugs. It is commonly used as spice, which is found in almost every kitchen to add flavour in many dishes especially in North India where it is used as an ingredient of garam masala. It is used in bakery and confectionary products, pudding, meats, sausages, sauces. In traditional systems of medicine such as Ayurveda and folk system this plant is used in treating variety of human ailments due to the presence of different kind of phytochemical constituents. It contains various essential oil constituents such as myristicin, elemicin, safrole, terpenes, alpha-pinene, beta-pinene, myristic acid and also present triacylglycerols like trimyristin, phenolic compounds, flavonoids, tocopherol, ascorbic acid, and many more other constituents are presents. The seeds and fruits parts of these plant are used in traditional practices for the treating various diseases like dysentery, weakness, sexual disorders, and darkness of complexion. The most common uses of *M. fragrans* is used as a carminative, anti-oxidant, analgesic, hypolipidemic, anti-pyretic, aphrodisiac, antiulcerogenic, anti-inflammatory, diuretic, hypnotic, antispasmodic, and stimulant. It is also used for treating the problems related to spleen, liver, and enhances the blood circulation, boosts up the brain functioning and enhance digestion. For centuries Nutmeg has been used all over the world as a valuable spice. Nutmeg has also been used in traditional remedies for stomach and kidney disorders along with its use in flavour enhancer of foods and beverages.



Figure- Fruits of *Myristica fragrans*

TRADITIONAL USES

Myristica fragrans is commonly known as "Jatiphala" in Ayurveda. It is a valuable medicinal plant. It is used in many Ayurvedic practices for the treatment of numerous diseases. It significantly works on vitiated vatta and kapha. Its topical applications it is use an anti-inflammatory, analgesic skin disorders. It is used in treating cold and headache and useful in removing the foul smell of wound. It is used as appetizer, anti-helminthic, enhances digestion, increases the liver metabolism, diarrhoea, dysentery, irritable bowel syndrome, constipation, lose motion, cholera, cardiac

problems. anti-mucolytic agent. It is helpful against cold, cough, asthma, and carminative. seed powder or decoction of the seeds used against diarrhoea and rheumatism. Chinese people use seed powder in the treatment of dysentery, digestive problem, piles, and leucorrhoea. The fruits of *M. fragrans* plant as steam-bathing material. They also use this plant in the treatment of rheumatism, syphilis, and toothache. The fruit powder of *M. fragrans* as a remedy for burning, general weakness, headache and insomnia people use seeds along with milk to enhance their digestion. The fruit of this plant is used in cooking as a spice and to enhance the flavour. The use of Jaiphal along with ajwain seeds and choti elaichi to treat diarrhoea among infants.

2. METHODOLOGY

PHARMACOLOGICAL ACTIVITIES

Antioxidant activity

Anti-oxidant Dorman et al., conducted an in-vitro study to check the anti-oxidant potential of many medicinal herbs against synthetic anti-oxidants like BHA (butylated hydroxy anisole), BHT (butylated hydroxytoluene), alpha tocopherol and pyrogallol. It was found from the study that essential oil extracted from nutmeg is an effective anti-oxidant agent.

Nutmeg possesses antioxidant activity due to the presence of various compounds including β caryophyllene and eugenol, having hydrogen atoms in the allylic or benzylic positions. Because of the comparatively simple abstraction of atomic hydrogen from these functional groups, these compounds have high antioxidant activity. The abstraction of atomic hydrogen is done by peroxy radicals that produced under oxidative stress. In another view, role of Eugenol in nutmeg favors the antioxidant property by promoting the activities of superoxide dismutase, catalase, glucose-6-phosphate dehydrogenase, glutathione peroxidase and glutamine transferase enzymes. The compounds having catechol like structure as in caffeic acid are considered to be good antioxidants as they easily donate electrons or phenolic hydrogen to the acceptors, such as lipid peroxyl groups or reactive oxygen species. Calliste et al., (2010) stated that lignan derivatives are considered as a class of compounds that shows the antioxidant potential of nutmeg seeds. After absorption of lignans and their glycosides into the body, they are metabolized to produce biologically active compounds having catechol structures that are responsible of high antioxidant property of nutmeg seeds.

Immuno-modulatory and radio-protective activities

The lignans present in fresh nutmeg and mace show radio modifying and immune modulatory properties, present in the aqueous extract of fresh nutmeg mace These properties found in cell free systems and protected PUC18 plasmid against radiation that induced DNA damage. The mammalian splenocytes in response to polyclonal T cell mitogen concanavalin A (Con A) proliferate. This process is inhibited by these mace lignans which was due to G1 phase of cell cycle and augmentation of apoptosis as presented by increase in pre G1 cells. The increase in activation of induced cell death by mace lignans was depending on the dosage. Splenocytes are protected by mace lignans against radiations. These radiations induced by producing intracellular reactive oxygen species depending on the dose. Mace lignans was not cytotoxic for lymphocytes. On the other hand, in splenocytes the radiation-induced DNA damage is inhibited by decreasing DNA fragmentation.

Antimicrobial activity

Anti-microbial Orabi et al., conducted an experimental study in 1991 to check the anti-microbial potential of mace of *M. fragrans* against *Staphylococcus aureus* and *Candida albicans*. It was revealed that resorcinols malabaricon B [1] and malabaricon C [2] both showed significant anti-microbial potential against *Staphylococcus aureus* and *Candida albicans*. As per the results of study conducted by Hattori et al., dehydro-di-isoeugenol and 5-methoxydehydrodiisogenol were found to be effective against *Streptococcus mutans*. They significantly helped in the inhibition of *Streptococcus mutans* growth.

The essential oil and different extracts of aromatic plants have shown strong antimicrobial activity against variety of fungi as well as bacteria. Narasimhan et al (2006) demonstrated the antibacterial activity by preparing chloroform extract of nutmeg against both gram negative and gram-positive bacteria. They found myristic acid and trimyristin are the main antibacterial compounds extracted from nutmeg seeds. Researchers isolated three lignans (mesodihydroguaiaretic acid, nectandrin-B and erythroaustrobailignan-6) showing antifungal activity, from the methanolic extract of nutmeg seeds. The development of wheat leaf rust and rice blast was suppressed by these three lignans. Some compounds like carvacrol, γ -cymene, α -pinene, β -pinene, and β caryophyllene are reported to be antimicrobial present in essential oil of nutmeg seeds. Some plant phenolics are also reported for antimicrobial activity.

Antifungal and anti-inflammatory activities

Antifungal and anti-inflammatory activities of plant essential oil are due to the presence of β -caryophyllene. α -Pinene and β -pinene which are monoterpene hydrocarbons are also antimicrobial agents. They are involved in membrane disruption. Carvacol is another significant compound for antimicrobial activity. Carvacol works in the same way as the other phenolic compounds, which work through membrane destruction, resulting in an increase in permeability of membrane to potassium ions and protons, proton-motive force disruption and intracellular ATP pool depletion. γ Cymene (a precursor of carvacrol) could also be an important component. It has been demonstrated that γ cymene works synergistically with carvacrol in membrane enlarging, which results weakening of the membrane while alone shows weak antibacterial activity. It has been proposed that antimicrobial activity is due to the minor and major both compounds; while it is feasible that the major compound controlled by other minor compounds.

Anti-inflammatory activity

Several authors reported anti-inflammatory activity of nutmeg as well as its oil. Like nonsteroidal anti-inflammatory drugs, pharmacological activities also exhibited by nutmeg oil. But anti-inflammatory activity is shown only by petroleum ether extracts. The total extract of nutmeg activated an enzyme that is AMP-activated protein kinase enzyme (potential therapeutic target) for curing the metabolic syndrome including type-2 diabetes and obesity. Seven compounds like tetrahydrofuroguaiacin B, 2,5-bis-aryl-3,4-dimethyltetrahydrofuranlignans, fragransin C1, saucernetindiol, nectandrin B, verrucosin, galbacin and nectandrin A were isolated from this extract as an active constituent. Some of the isolated compounds produced strong AMPK stimulation in differentiated C2C12 cells, at 5 μ M concentration. Nutmeg and its active components not only used to treat type-2 diabetes and obesity but also for the development of agent's other metabolic disorders.

Anti-carcinogenic and hepatoprotective activity-

Anti-cancer Prakash et al., studied the anti-cancerous behavior of *Myristica fragrans* in an in-vitro study on human cell lines Colon cell (Colon502713, Colo205), Liver (Hep-2), Lung (A-549), Ovary OVCAR-5 and Prostrate (PC-5) and central nervous system. Ethanolic extract of seed of *Myristica fragrans* (Houtt) was found to be effective against these cell lines. Ethanolic extract exhibited highest anti-cancerous activity against OVCAR-5. The results of this study demonstrated that *Myristica fragrans* is a potent anti-cancer agent.

Nutmeg shows resistance against carcinogenic elements. Hussain and Rao, (1991) reported that, in Swiss albino mice uterine cervix, 3-methylcholanthrene -induced carcinogenesis could be prohibited by mace oral administration. Nutmeg also shows hepatoprotective activity. This property observed in rats with damaged liver, by giving nutmeg in their diets. Kyriakis et al., (1994) studied on the activities of hepatic carcinogen-metabolizing enzymes, like aryl hydrocarbon hydroxylase, cytochrome P450, and acid soluble sulphhydryl and glutathione-Transferase level in albino mice and checked the influence of essential oil from nutmeg. They found that the essential oil hinders the activities of the host enzymes related with detoxication and activation of xenobiotic components, as well as mutagens and chemical carcinogens.

Hepatoprotective Zhao et al., conducted an experimental study to check the effects of AEN (alcohol extract of nutmeg, the seed of *Myristica fragrans* on obesity and inflammation which are associated with Non-alcoholic fatty liver disease (NAFLD). They use mice models for the study. Models were fed with high-fat-diet. It was revealed from the study, alcohol extract of nutmeg significantly reduced the body weight and levels of inflammation, cholesterol and lipid accumulation, blood glucose which supports its use as an hepatoprotective agent. Dkhil et al., also studied hepatoprotective activity of *Myristica fragrans*. The study was carried out on rat models. Models were administered with paracetamol (acetaminophen; N-acetyl-p-amino-phenol (APAP) for the artificial induction of hepatotoxicity. *Myristica fragrans* Kernel extract showed some significant effect against APAP induced toxication.

Aphrodisiac Activity-

Ahmed et al., conducted a study on male rat models to evaluate the activity of *Myristica fragrans* Houtt. (nutmeg) on sexual behavior. It was observed that 50% ethanolic extract of the plant exhibited aphrodisiac activity.

Anti-stress activity-

was examined by Dhingra et al., on mice models. Stress was induced in the models by using two methods that are the forced swim test (FST) and the tail suspension test (TST) method. It was observed that *M. fragrans* extract significantly reduced the immobility periods of mice in both the FST and the TST methods.

Anti-diarrhoeal -

A study was conducted by Gupta et al., on ileal loops of rabbit and guinea pig. E. coli heat labile (LT) and heat stable (ST) enterotoxins were used in this experimental study. The result revealed that hexane-soluble fraction was

associated with inhibition of secretory responses which were induced by *E. coli* heat labile (LT) and heat stable (ST) enterotoxins.

Antithrombotic-

As per the data report of an in-vitro study carried out by Janssens et al., it was revealed that eugenol and iso-eugenol were associated with inhibition of platelet aggregation, this attributes to its antithrombotic activity.

Hypolipidemic effect-

In-vivo study conducted on rabbit models by Ram et al., suggested that the oral administration of ethanolic extract of nutmeg at the dosage of 500 mg/kg at the regular interval of 60 days in artificially induced hypolipidemic rabbits significantly lowered the lipoprotein lipids levels which supports its use as an hypolipidemic agent. Sharma et al., carried out an in-vivo study on hypercholesterolemic rabbit models which revealed that administration of seed extract of *M. fragrans* exhibited properties like reduction in serum cholesterol and LDL cholesterol by 69.1 and 76.3%. It also lowered down the ratio of cholesterol/phospholipid by 31.2% and it significantly elevated the ratio of decreased HDL.

Anti-diabetic -

As per the study report of Han et al., macelignan extracted from *M. fragrans* is associated with anti-diabetic potential. It activates the peroxisome proliferator receptor (PPAR, α/δ) which helps in improving the insulin sensitivity and lipid metabolic disorders.

Anti-convulsant-

Wahab et al., studied anti-convulsant potential of *M. fragrans* in well-established animal seizure models. It was observed that nutmeg oil exhibits anti-convulsant properties. During the study nutmeg oil showed anti-convulsant activity against tonic seizures induced by pentylenetetrazole which supports its use as an anticonvulsant agent.

3. CONCLUSION

M. fragrans is a mediational herb which is most used in traditional systems of medicine for treating various human ailments. It is widely used for the treatment of several diseases Apart from its therapeutic uses, it is well known spice used in almost all type of cooking. *Myristica fragrans* is a resource of medicinally active compounds and has diverse pharmacological effects; hence, this drug encourages researchers to explore its various novel therapeutic uses for the benefit of mankind.

4. REFERENCE

- [1] NAIR, R., KALARIYA, T., & Chanda, S. Antibacterial activity of some selected Indian medicinal flora. Turkish Journal of biology, (2005); 29(1): 41-47.
- [2] Ghorbani, A. Studies on pharmaceutical ethnobotany in the region of Turkmen Sahra, north of Iran:(Part 1): General results. Journal of ethnopharmacology, (2005); 102(1): 58-68.
- [3] Chukwuma, E. C., Soladoye, M. O., & Feyisola, R. T. Traditional medicine and the future of medicinal Plants in Nigeria. Journal of Medicinal Plants Studies, (2015); 3(4) : 23-29.
- [4] Kirtikar KR, Basu BD: Edited by E. Blatter. Indian Medicinal Plants. 2nd ed. Vol III, Deharadun;1984, PG 2141
- [5] Prof. LucasDS. DravyaGunaVigyanaStudy of DravyaMateriaMedica, 1st ed. Vol II, Varanasi; ChaukhambhaVishwabharati: 369, (2008).
- [6] The Ayurvedic Pharmacopoeia of India. 1sted. Part 1, Vol 2, New Delhi; Government of India, Ministry of Health and Family Welfare, Department of ISM&H: p.68.(1999).
- [7] Chopra, ChopraIC, HandaKL.Kapur LD. Indigenous drugs of India, 2nd reprint, Calcutta; Academic publishers: 200, (1994).
- [8] Garg A, Kumar A, Nair A, Reddy A. Analysis of some Indian medicinal herbs by INAA. Journal of radioanalytical and nuclear chemistry, (2007); 271(3): 611-9.
- [9] Baldi A. Erectile Dysfunction and Its Treatment.
- [10] Sharma PC, Yelne MB, Dennis TJ, Joshi A, Billore KV. Database on medicinal plants used in Ayurveda, (2002); 5: 213-217.
- [11] Eshwara JH, Kulathunga RD, Gunarathna ED. (Standardization of Nagaraja Guliya; A Sri Lankan Traditional Formula Used in Poisons of Animal Origin (Jangama Visha). Asian Journal of Advanced Research and Reports, (2020); 19: 49-59.
- [12] Muddgal D. Dravyagun Vijnana. Ayurvedic hindi pustak bhandar.
- [13] Sharma PV. Dravyagun Vigyan. Chaukhambha Bharti Academy, Varanasi, (2019).

- [14] Desai A, Desai C, Desai H, Mansuri A, Desai J. Possible Role of Medicinal Plants in COVID19-a Brief Review. *International Journal of Scientific Development and Research*, (2020); 5(4): 205-9.
- [15] Tambekar DH, Dahikar SB. Screening of Some Herbal Preparations Used In Indian Ayurvedic Medicine For Its Antibacterial Potentials. *International Journal of Pharmaceutical Sciences and Research*, (2011); 2(2):
- [16] Russo E. Cannabis in India: ancient lore and modern medicine. In *Cannabinoids as therapeutics*, (2005); 1-22.
- [17] Van Gils C, Cox PA. Ethnobotany of nutmeg in the Spice Islands. *Journal of ethnopharmacology*, (1994); 42(2): 117-24.
- [18] Sihotang, V. B. L., Yang, G., Chi, X., & Huang, L. The Ethnobotanical Survey of Clove, Pepper, and Nutmeg and Their Utilization by Chinese and Indonesian People. *Journal of Tropical Biology & Conservation (JTBC)*, (2018); 15â-27.
- [19] A. A., & Bello, O. A. Ethnobotanical potentials of common herbs in Nigeria: A case study of Enugu state. *Educational Research and Reviews*, (2006); 1(1): 16-22.
- [20] Kayode, J., & Ogunleye, T. O. Checklist and status of plant species used as spices in Kaduna State of Nigeria. *Research Journal of Botany*, (2008); 3(1): 35- 40.
- [21] Suwardi AB, Navia ZI, Harmawan T, Mukhtar E. Ethnobotany, and conservation of indigenous edible fruit plants in South Aceh, Indonesia. *Biodiversitas Journal of Biological Diversity*, (2020); 21(5).
- [22] Wu M, Guo P, Tsui SW, Chen H, Zhao Z. An ethnobotanical survey of medicinal spices used in Chinese hotpot. *Food research international*, (2012); 48(1): 226- 32.
- [23] Mitra R, Mitchell B, Gray C, Orbell J, Coulepis T, Muralitharan M. Medicinal plants of Indonesia. *Asia Pacific biotech news*, (2007); 11(11): 726-43.
- [24] Ouhaddou H, Boubaker H, Msanda F, El Mousadik A. An ethnobotanical study of medicinal plants of the Agadir Ida Ou Tanane province (southwest Morocco). *Journal of Applied Biosciences*, (2014); 84: 7707-22.
- [25] Jianwitchayakul P, Aung SS, Aye T, Zarli M, Myint ZM, Mu AA. Investigation Biodiversity Ethnobotany in Mon State, Republic of the Union of Myanmar, (2018); 11-20.
- [26] Dorman, H. D., Surai, P., & Deans, S. G. In vitro antioxidant activity of a number of plant essential oils and phytoconstituents. *Journal of Essential Oil Research*, (2000); 12(2): 241-248.
- [27] Jaiswal, P., Kumar, P., Singh, V. K., & Singh, D. K. Biological effects of *Myristica fragrans*. *Annual review of biomedical sciences*, (2009); 11: 21-29.
- [28] Shahzadi, I., Nadeem, R., Hanif, M. A., Mumtaz, S., Jilani, M. I., & Nisar, S. Chemistry and biosynthesis pathways of plant oleoresins: important drug sources. *IJCBS*, (2017); 12: 18-52.
- [29] Widelski, J., & Kukula-Koch, W. A. Psychoactive Drugs. In *Pharmacognosy* (pp. 363- 374). Academic Press, (2017).
- [30] Orabi, K. Y., Mossa, J. S., & El-Feraly, F. S. Isolation and characterization of two antimicrobial agents from mace (*Myristica fragrans*). *Journal of natural products*, (1991); 54(3): 856-859.
- [31] Hattori M, Hada S, Watahiki A, Ihara H, Shu YZ, Kakiuchi N, Mizuno T, Namba T. Studies on dental caries prevention by traditional medicines. X.: antibacterial action of phenolic components from mace against *Streptococcus mutans*. *Chemical and pharmaceutical bulletin*, (1986); 34(9): 3885-93.
- [32] Calliste, C. A., Kozłowski, D., Duroux, J. L., Champavier, Y., Chulia, A. J., & Trouillas, P. A new antioxidant from wild nutmeg. *Food chemistry*, (2010); 118(3): 489-496.
- [33] Checker, R., Chatterjee, S., Sharma, D., Gupta, S., Variyar, P., Sharma, A., & Poduval, T. B. Immunomodulatory and radioprotective effects of lignans derived from fresh nutmeg mace (*Myristica fragrans*) in mammalian splenocytes. *International immunopharmacology*, (2008); 8(5): 661-669.
- [34] Hanif, M. A., Bhatti, H. N., Jamil, M. S., Anjum, R. S., Jamil, A., & Khan, M. M. Antibacterial and antifungal activities of essential oils extracted from medicinal plants using CO2 supercritical fluid extraction technology. *Asian journal of chemistry*, (2010); 22(10): 7787.
- [35] Cho, J. Y., Choi, G. J., Son, S. W., Jang, K. S., Lim, H. K., Lee, S. O., ... & Kim, J. C. Isolation, and antifungal activity of lignans from *Myristica fragrans* against various plant pathogenic fungi. *Pest Management Science: formerly Pesticide Science*, (2007); 63(9): 935-940.
- [36] Dorman, H. D., & Deans, S. G. Chemical composition, antimicrobial and in vitro antioxidant properties of *Monarda citriodora* var. *citriodora*, *Myristica fragrans*, *Origanum vulgare* ssp. *hirtum*, *Pelargonium* sp. and *Thymus zygis* oils. *Journal of Essential Oil Research*, (2004); 16(2) : 145-150.
- [37] Sabulal, B., Dan, M., Kurup, R., Pradeep, N. S., Valsamma, R. K., & George, V. Caryophyllene-rich rhizome oil of *Zingiber nimmonii* from South India: Chemical characterization and antimicrobial activity. *Phytochemistry*, (2006); 67(22): 2469-2473.

- [38] Dorman, H. D., & Deans, S. G. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *Journal of applied microbiology*, (2000); 88(2): 308-316.
- [39] Hussain, S. P., & Rao, A. R. Chemopreventive action of mace (*Myristica fragrans*, Houtt) on methylcholanthrene-induced carcinogenesis in the uterine cervix in mice. *Cancer letters*, (1991); 56(3): 231-234.
- [40] Kyriakis, J. M., Banerjee, P., Nikolakaki, E., Dai, T., Rubie, E. A., Ahmad, M. F., ... & Woodgett, J. R. The stress-activated protein kinase subfamily of c-Jun kinases. *Nature*, (1994); 369(6476): 156-160.
- [41] Mueller, M., Hobiger, S., & Jungbauer, A. Anti-inflammatory activity of extracts from fruits, herbs and spices. *Food chemistry*, (2010); 122(4): 987-996.
- [42] Olajide, O. A., Makinde, J. M., & Awe, S. O. Evaluation of the pharmacological properties of nutmeg oil in rats and mice. *Pharmaceutical biology*, (2000); 38(5): 385-390.
- [43] Nguyen, P. H., Le, T. V. T., Kang, H. W., Chae, J., Kim, S. K., Kwon, K. I., ... & Oh, W. K. AMP-activated protein kinase (AMPK) activators from *Myristica fragrans* (nutmeg) and their antiobesity effect. *Bioorganic & medicinal chemistry letters*, (2010); 20(14): 4128-4131.
- [44] Prakash, E., & Gupta, D. K. Cytotoxic activity of ethanolic extract of *Myristica fragrans* (Houtt) against seven human cancer cell lines. *Journal of Food and Nutrition Science*, (2013); 1(1): 1-3.
- [45] Ultee, A., Bennik, M. H. J., & Moezelaar, R. J. A. E. M. The phenolic hydroxyl group of carvacrol is essential for action against the foodborne pathogen *Bacillus cereus*. *Applied and environmental microbiology*, (2002); 68(4): 1561-1568.
- [46] Bounatirou, S., Smiti, S., Miguel, M. G., Faleiro, L., Rejeb, M. N., Neffati, M., ... & Pedro, L. G. Chemical composition, antioxidant and antibacterial activities of the essential oils isolated from Tunisian *Thymus capitatus* Hoff. et Link. *Food chemistry*, (2007); 105(1): 146-155.
- [47] Zhao, W., Song, F., Hu, D., Chen, H., Zhai, Q., Lu, W., ... & Wang, G. The protective effect of *Myristica fragrans* Houtt. Extracts against obesity and inflammation by regulating free fatty acids metabolism in nonalcoholic fatty liver disease. *Nutrients*, (2020); 12(9): 2507.
- [48] Dkhil, M. A., Abdel Moneim, A. E., Hafez, T. A., Mubarak, M. A., Mohamed, W. F., Thagfan, F. A., & Al-Quraishy, S. *Myristica fragrans* kernels prevent paracetamol-induced hepatotoxicity by inducing anti-apoptotic genes and Nrf2/HO-1 pathway. *International journal of molecular sciences*, (2019); 20(4): 993.
- [49] Ahmad, S., Latif, A., Qasmi, I. A., & Amin, K. M. Y. An experimental study of sexual function improving effect of *Myristica fragrans* Houtt. (nutmeg). *BMC Complementary and Alternative Medicine*, (2005); 5(1): 1-7.
- [50] Parle, M., Dhingra, D., & Kulkarni, S. K. Improvement of mouse memory by *Myristica fragrans* seeds. *Journal of medicinal food*, (2004); 7(2): 157-161.
- [51] Dhingra, D., & Sharma, A. Antidepressantlike activity of n-hexane extract of nutmeg (*Myristica fragrans*) seeds in mice. *Journal of medicinal food*, (2006); 9(1): 84-89.
- [52] Gupta, S., Yadava, J. N. S., Mehrotra, R., & Tandon, J. S. Anti-diarrhoeal profile of an extract and some fractions from *Myristica fragrans* (Nutmeg) on *Escherichia coli* enterotoxin-induced secretory response. *International journal of pharmacognosy*, (1992); 30(3): 179-183.
- [53] Janssens, J., Laekeman, G. M., Pieters, L. A., Totte, J., Herman, A. G., & Vlietinck, A. J. Nutmeg oil: identification and quantitation of its most active constituents as inhibitors of platelet aggregation. *Journal of ethnopharmacology*, (1990); 29(2): 179-188.
- [54] Ram, A., Lauria, P., Gupta, R., & Sharma, V. N. Hypolipidaemic effect of *Myristica fragrans* fruit extract in rabbits. *Journal of ethnopharmacology*, (1996); 55(1): 49-53.
- [55] Sharma, A. R. T. I., Mathur, R. I. T. U., & Dixit, V. P. Prevention of hypercholesterolemia and atherosclerosis in rabbits after supplementation of *Myristica fragrans* seed extract. *Indian journal of physiology and pharmacology*, (1995); 39: 407-410.
- [56] Han, K. L., Choi, J. S., Lee, J. Y., Song, J., Joe, M. K., Jung, M. H., & Hwang, J. K. Therapeutic potential of peroxisome proliferators-activated receptor α/γ dual agonist with alleviation of endoplasmic reticulum stress for the treatment of diabetes. *Diabetes*, (2008); 57(3): 737-745.
- [57] Wahab, A., Haq, R. U., Ahmed, A., Khan, R. A., & Raza, M. Anticonvulsant activities of nutmeg oil of *Myristica fragrans*. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, (2009); 23(2): 153-158.