

ANTIDIABETIC & ANTIOXIDANT ACTIVITY ON THE STUDY OF BUTTERFLY PEA LEAF EXTRACT

Ms. Roshanee D Agrawal¹, Mr. Mahesh K. Patil², Ms. Sonali B. Jagdhane³,

Mr. Ajit G. Jadhav⁴, Mr. Avinash P. Deore⁵

^{1,2,3,4,5}Aditya Institute of Pharmacy, Chalisgaon, India.

ABSTRACT

Clitoria ternatea, often known as butterfly pea plant, is widely utilized in Ayurvedic and traditional medicine. It is also utilized in the culinary and cosmetic industries for a variety of applications. This study examined the antibacterial and anticancer properties of methanol leaf extract from Clitoria ternatea against human cells. Promyelocytic Leukemia Cells Phytochemical screening identifies secondary metabolites in the extract, including alkaloids, phlobatannins, triterpenoids, flavonoids, lipids, steroids, terpenoids, tannins, and glycosides. Leaf extract included ether and carboxylic acid groups. The plant has numerous pharmacological benefits, including antioxidant, hypolipidemic, anticancer, anti-inflammatory, analgesic, antipyretic, antidiabetic, CNS, antibacterial, gastro-intestinal, antiparasitic, and insecticidal properties.

Key words- CNS, antibacterial, gastro-intestinal, antiparasitic, and insecticidal properties.

1. INTRODUCTIONS

Asian Pigeonwings Flower (Clitoria ternatea) is a popular herbal plant used in traditional medicine. A som of the Pigeonwings flower has active elements that make it an antibacterial chemical. The active compounds include alkaloids, phenols, saponins, tannins, and flavonoids. Each of these active chemicals has a unique Mechanism for limiting bacterial growth. Many patients worldwide use medicinal plants and herbs to improve their health. Conducting scientific research on their medicinal potential, biological features, and safety can help make informed judgments regarding their use There are hundred so fimportant medications, and biologically Active chemicals were derived from traditional medicinal herbs. The plant has numerous pharmacological actions, such as antibacterial, antioxidant, anticancer, hypolipidemic, cardiovascular, central nervous, respiratory, immunological, anti-inflammatory, analgesic, and antipyretic. It has several pharmacological actions, including anti-inflammatory, analgesic, antipyretic, antidiabetic, CNS, antibacterial, gastro-intestinal antiparasitic, and insecticidal properties. This review will focus on the chemical composition and pharmacological effects of Clitoria ternatea.



Plant biology-

Synonyms:

Clitoria albiflora Mattei, *Clitoria bracteata* Poir., *Clitoria mearnsii* De Wild., *Clitoria tanganicensis* Micheli,
Clitoria zanzibarensis Vatke

Taxonomic classification:

Kingdom: Plantae

Subkingdom: Viridaeplanta

Infrakingdom: Streptophyta;

Division: Tracheophyta

Sub-family: Papilionaceae

Class: Magnoliopsida;

Superorder: Rosanae

Order: Fabales;

Family: Leguminosae

Genus: Clitoria L.

Species: *Clitoria ternatea*

Common name - Butterfly pea, Clitoria ternatea, Asian Pigeon wings flower, Blue pea

Description-

Leaflets ovate to elliptic-oblong, up to 6.5×4 cm, mostly hairless above,

Pubescent below. Flowers axillary, solitary or 2 together, resupinate, large and showy, bright blue

Collection and cultivation-


Clitoria ternatea leaves were obtained from south America & Asia . Using whose blue flowers are commonly used as a food dye, the leaves collected and converted into powder form.

Chemical constituents:

The plant has a variety of chemical elements, including tannins, phlobatannin, polysaccharides, saponins, triterpenoids, phenols, flavonoids, alkaloids.

Anthraquinone, anthocyanins, cardiac glycosides, Stigmast-4-ene-3,6-dione, volatile oils, and steroids Clitoria ternatea seeds include fatty acids such as palmitic, stearic, oleic, linoleic, and linolenica.

Extraction process of Asian pigeon wings



Pharmacological effects-

Antimicrobial effect:

The antibacterial properties of Clitoria ternatea were studied using agar disc and well diffusion methods. Clitoria ternatea leaf extracts in organic solvents (petroleum ether, ethyl acetate, and methanol) were evaluated against *Bacillus cereus*, *Staphylococcus aureus*, *Klebsiella pneumonia*, *Proteus vulgaris*, and *Salmonella. typhi*. The results indicated good antibacterial efficacy against the investigated microbiological pathogens. Methanol extract was shown to have higher inhibitory action than petroleum ether and ethyl acetate.

Anti-inflammatory antipyretic analgesic effects:

The petroleum ether extract ($60-80^{\circ}\text{C}$) had substantial anti-inflammatory and analgesic effects. The methanolic extract of Clitoria ternatea Linn. leaves was tested for analgesic activity on mice at doses of 200 and 400 mg/kg body weight. The acetic acid-induced writhing test was used to assess analgesic activity.

Anticancer effect:

The in vitro cytotoxic effect of petroleum ether and ethanolic floral extracts (10, 50, 100, 200, 500 $\mu\text{g/ml}$) of Clitoria ternatea was examined using the trypan blue dye exclusion method. Both extracts showed considerable dose-dependent cell cytotoxic action. At a concentration of 10 $\mu\text{g/ml}$, petroleum ether extract reduced cell count by 8%.

However, at 500 $\mu\text{g}/\text{ml}$, the reduction was 100%. The ethanolic extract reduced cell count by 1.33% at a dosage of 10 $\mu\text{g}/\text{ml}$ and by 80% at 500 $\mu\text{g}/\text{ml}$. The cytotoxicity of *Clitoria ternatea* flower extracts (aqueous and methanol) was tested on six cell lines, including normal and cancerous. These included the hormone-dependent breast cancer cells.

Central nervous effect:

Clitoria ternatea seeds and leaves are traditionally used as a brain tonic to improve memory and intelligence. *Clitoria ternatea* was researched in Alzheimer's disease to determine its efficacy and identify the main bioactive ingredient responsible for its activity. Ayurveda is widely utilized for several CNS effects, including memory augmentation. The origin of shankhpushpi is unclear due to the use of various plants in different parts of India.

Benefits



1. Improve cognitive performance and reduce symptoms of common diseases fever, inflammation.
2. Butterfly pea flower tea, often known as blue tea, is prepared from the leaf.
3. Butterfly pea flower, cultivated widely in Southeast Asia, produces a vibrant tea.
4. The flavor is flowery and gently sweet, reminiscent of chamomile
5. Blue tea contains anthocyanins, which can help manage blood sugar levels.
6. Butterfly pea blossoms are a natural alternative to artificial blue food colorants, which are becoming less popular due to health concerns.

2. RESULTS AND DISCUSSION

The maceration extraction procedure involves sample immersion, which might Cell wall rupture occurs when concentrations inside and outside the cell differ, attracting chemicals in the cytoplasm to the solvent.

3. CONCLUSION

Clitoria ternatea is a potential medicinal plant with a wide spectrum of pharmacological properties. Its efficiency and safety make it suitable for various medical uses.

4. REFERENCE

- [1] Al-Snafi AE. Therapeutic properties of medicinal plants: a review of their effect on reproductive systems (part 1). *Ind J of Pharm Sci & Res* 2015; 5(4): 240-248.
- [2] Al-Snafi AE. Therapeutic properties of medicinal plants: a review of their gastro-intestinal effects (part 1). *Ind J of Pharm Sci & Res* 2015; 5(4): 220-232.
- [3] Al-Snafi AE. Therapeutic properties of medicinal plants: a review of their antiparasitic, antiprotozoal, molluscicidal and insecticidal activity (part 1). *J of Pharmaceutical Biology* 2015; 5(3): 203-217.
- [4] Al-Snafi AE. Therapeutic properties of medicinal plants: a review of plants with antidiabetic effects (part 1). *J of Pharmaceutical Biology* 2015; 5(3): 218-229.
- [5] Syarifuddin, A., N. Sulistyani, dan Kintoko. Profil KLT-Bioautografi dan Densitometri Fraks Teraktif (IsolatKp13) dari Bakteri Rizosfer Kayu Putih (*Melaleuca leucadendron* L.). *Jurnal Farmasi Sains dan Praktis*. 2019;5(1):27-33.
- [6] Kurama, G.M., W. Maarisit, E.Z. Karundeng, dan N.O. Patalangi. Uji Aktivitas Antibakteri Ekstrak Etanol Daun Benalu Langsat (*Dendrophoe* sp.) Terhadap Bakteri *Klebsiella pneumoniae*. *Jurnal Biofarmasetikal Tropis*. 2020;3(2):27-33.
- [7] Katuuk, R.H.H., S.A. Wanget, dan P. Tumewu. Pengaruh Perbedaan Ketinggian Tempat terhadap Kandungan Metabolit Sekunder pada Gulma Babadotan (*Ageratum conzyoides* L.). *COCOS*. 2019.
- [8] Jacob L, Latha MS. Anticancer activity of *Clitoria ternatea* Linn. against Dalton's lymphoma. *International Journal of Pharmacognosy and Phytochemical Research*. 2012; 4(4):207-12.
- [9] Jamil N, Zairi MNM, Nasim NAIM, Pa'ee F. Influences of environmental conditions to phytoconstituents in *Clitoria ternatea* (Butterfly Pea Flower)-A Review. *Journal of Science and Technology*. 2018; 10(2).

- [10] Kamilla L, Mansor SM, Ramanathan S, Sasidharan S. Effects of *Clitoria ternatea* leaf extract on growth and morphogenesis of *Aspergillus niger*. *Microscopy and Microanalysis*. 2009; 15(4):366–72. <https://doi.org/10.1017/S1431927609090783>. PMid:19575837
- [11] Kumar D, Dhobi M. Antianxiety and antioxidant profile of blue and white variety of *Clitoria ternatea* L. *Indian Journal of Research in Pharmacy and Biotechnology*. 2016; 4(3):90.
- [12] Lijon MB, Meghla NS, Jahedi E, Rahman MA, Hossain I. Phytochemistry and pharmacological activities of *Clitoria ternatea*. *International Journal of Natural and Social Sciences*. 2017; 4(1):1–10.
- [13] Mathew N, Anitha MG, Bala TSL, Sivakumar SM, Narmadha R, Kalyanasundaram M. Larvicidal activity of *Saraca indica*, *Nyctanthes arbor-tristis*, and *Clitoria ternatea* extracts against three mosquito vector species. *Parasitology Research*. 2009; 104(5):1017–25. <https://doi.org/10.1007/s00436-008-1284-x>. PMid:19039604
- [14] Nguyen KNT, Nguyen GKT, Nguyen PQT, Ang KH, Dedon PC, Tam JP. Immunostimulating and Gram-negative-specific antibacterial cyclotides from the butterfly pea (*Clitoria ternatea*). *The FEBS Journal*. 2016; 283(11):2067–90. <https://doi.org/10.1111/febs.13720>. PMid:27007913
- [15] Ramaswamy V, Varghese N and Simon A. An investigation on cytotoxic and antioxidant properties of *Clitoria ternatea* L. *International Journal of Drug Discovery* 2011; 3(1): 74-77.
- [16] Jacob L and Latha MS. Anticancer activity of *Clitoria ternatea* Linn. against Dalton's lymphoma. *International Journal of Pharmacognosy and Phytochemical Research* 2012; 4(4); 207-212.
- [17] 16. Mukhopadhyay R, Bhattacharya S and Biswas M. In vitro free radical scavenging activity of *Clitoria ternatea* leaf extracts. *Journal of Advanced Pharmacy Education & Research* 2012; 2(4):206-209.
- [18] Kamilla L, Mansor SM, Ramanathan S and Sasidharan S. Effects of *Clitoria ternatea* leaf extract on growth and morphogenesis of *Aspergillus niger*. *Microsc Microanal* 2009; 15(4) :366-372.
- [19] 18. Kelemu S, Cardona C and Segura G. Antimicrobial and insecticidal protein isolated from seeds of *Clitoria ternatea*, a tropical forage legume. *Plant Physiol Biochem* 2004;42(11):867-873.
- [20] Shekhawat N and Vijayvergia R. Anthelmintic activity of extracts of some medicinal plants. *International Journal of Computational Science and Mathematics* 2011;3(2):183-187.
- [21] Nirmal SA, Bhalke RD, Jadhav RS and Tambe VD. Anthelmintic activity of *Clitoria ternatea*. *Pharmacologyonline* 2008; 1: 114-119.