

www.ijprems.com

editor@ijprems.com

# INTERNATIONAL JOURNAL OF PROGRESSIVE RESEARCH IN ENGINEERING MANAGEMENT AND SCIENCE (IJPREMS)

e-ISSN : 2583-1062

Impact Factor : 7.001

(Int Peer Reviewed Journal)

Vol. 05, Issue 02, February 2025, pp : 1199-1201

# SYNTHESIS AND CHARACTERIZATION OF MIXED LIGAND COMPLEXES OF 1-{(E)-[(4-METHYL-2 NITROPHENYL) IMINO] METHYL} NAPHTHALEN-2-OL AND 2-{(E)-[(2,3-DIMETHYLPHENYL) IMINO]METHYL}PHENOL WITH CO(II), NI(II), CU(II) AND ZN(II) IONS.

Atmaram K. Mapari<sup>\*1</sup>

<sup>\*1</sup>Department Of Chemistry, Ramnarain Ruia Autonomous College, Matunga (E), Mumbai,

Maharashtra, India.

DOI: https://www.doi.org/10.58257/IJPREMS38741

# ABSTRACT

The mixed ligand complexes of Co(II), Ni(II), Cu(II) and Zn(II) with Schiff bases 1-{(E)-[(4-methyl-2-nitrophenyl)imino]methyl}naphthalen-2-ol (L<sup>1</sup>H) and 2-{(E)-[(2,3-dimethylphenyl)imino]methyl}phenol (L<sup>2</sup>H) have been synthesized and characterized. The resulting complexes were characterized by thermogravimetric analysis, magnetic moment measurements, conductivity measurements, IR, UV-visible spectral studies. The Schiff bases acts as bidentate monobasic ligands, coordinating through deprotonated phenolic oxygen and azomethine nitrogen atoms. The complexes are non-electrolytic in DMSO. The presence of the two coordinated water molecules in these complexes was indicated by IR spectra and thermogravimetric analysis of the complexes. From the analytical and spectral data the stoichiometry of these complexes have been found to be  $[M(L^1)(L^2)(H_2O)_2]$  {where M = Co(II), Ni(II), Cu(II) and Zn(II) complexes exhibited octahedral geometry.

Keywords: Schiff Bases, Mixed Ligand, Metal Complexes.

## **1. INTRODUCTION**

In the field of coordination chemistry, Schiff base metal complexes have a curious history [1]. The present work is the study of mixed ligand complexes of Co(II), Ni(II), Cu(II) and Zn(II) with Schiff bases 1-{(E)-[(4-methyl-2-nitrophenyl)imino]methyl}naphthalen-2-ol (L<sup>1</sup>H) and 2-{(E)-[(2,3-dimethylphenyl)imino]methyl}phenol (L<sup>2</sup>H). The structure of ligands L<sup>1</sup>H and L<sup>2</sup>H are shown in figure 1 and 2 respectively.



Figure 1: 1-{(E)-[(4-methyl-2-nitrophenyl)imino]methyl}naphthalen-2-ol

 $(L^1H)$ 



**Figure 2**: 2-{(E)-[(2,3-dimethylphenyl)imino]methyl}phenol (L<sup>2</sup>H)

# 2. MATERIALS AND METHODS

## Synthesis of ligands

The ligands ( $L^{1}H$  and  $L^{2}H$ ) were prepared by the condensation of the corresponding aldehyde and amine in 1:1 molar ratio by refluxing in ethanol [2].

#### **Preparation of complexes**

The mixed ligand metal complexes were prepared by precipitation method [3].

A4	<b>INTERNATIONAL JOURNAL OF</b>	e-ISSN :	
IJPREMS	PROGRESSIVE RESEARCH IN ENGINEERING	2583-1062	
	MANAGEMENT	<b>—</b>	
	AND SCIENCE (IJPREMS)	Impact	
www.ijprems.com	(Int Peer Reviewed Journal)	Factor : 7.001	
editor@ijprems.com	Vol. 05, Issue 02, February 2025, pp : 1199-1201		

# 3. RESULTS AND DISCUSSIONS

The resulting complexes were characterized by thermogravimetric analysis, magnetic moment measurements, IR and UV-visible studies.

#### **Conductivity Measurements**

The molar conductance values of Co(II), Ni(II), Cu(II) and Zn(II) complexes are 0.013, 0.018, 0.015 and 0.021 ohm<sup>-1</sup> mol<sup>-1</sup> cm<sup>2</sup> respectively of  $10^{-3}$  M solution in DMSO indicate that the metal complexes are non-electrolytic in nature.

### Electronic spectra and Magnetic moment

The electronic spectral studies of Mixed Ligand metal Complexes carried out in DMSO solution. The absorption spectrum of the Co(II) complex shows bands at ~ 10013 cm<sup>-1</sup> ( $\epsilon$  ~ 15) and ~ 19604 cm<sup>-1</sup> ( $\epsilon$  ~ 45) attributed to  ${}^{4}T_{1g}(F) \rightarrow {}^{4}T_{2g}(F)$  (v<sub>1</sub>) and  ${}^{4}T_{1g}(F) \rightarrow {}^{4}T_{1g}(P)$  (v<sub>3</sub>) transitions respectively in an octahedral field. The Co(II) complex has magnetic moment 5.14 BM also suggest an octahedral geometry [3], [4].. Ni(II) complex exhibits two electronic spectral bands at ~ 10613 cm<sup>-1</sup> ( $\epsilon$  ~ 20) and ~ 16564 cm<sup>-1</sup> ( $\epsilon$  ~ 61) which can be assigned to  ${}^{3}A_{2g}(F) \rightarrow {}^{3}T_{2g}(F)$  (v<sub>1</sub>) and  ${}^{3}A_{2g}(F) \rightarrow {}^{3}T_{1g}(F)$  (v<sub>2</sub>) transitions in an octahedral field<sup>5</sup>. The Ni(II) complex has magnetic moment 3.42 BM also suggest an octahedral geometry [3], [4]. The Cu(II) complex exhibit broad band centered at ~ 14704 cm<sup>-1</sup> ( $\epsilon$  ~ 77) mainly due to  ${}^{2}E_{g} \rightarrow {}^{2}T_{2g}$  transition suggesting the distorted octahedral geometry. The observed magnetic moment value for Cu(II) complex is 1.82 BM suggestive of distorted octahedral nature for the complex [3], [4]. Zn(II) complex does not exhibit any characteristic d-d transitions and is also found to be diamagnetic in nature [5].

### **Infrared** spectra

The important infrared frequencies exhibited by the ligands  $L^{1}H$  and  $L^{2}H$  and their mixed ligand complexes are given in the Table 1. Infrared spectra of the schiff bases  $L^{1}H$  and  $L^{2}H$  show a broad band centered at around 3453 and 3446 cm<sup>-1</sup> due to the phenolic hydroxyl group respectively in free ligands, which disappeared in spectra of their complexes indicating probably the coordination through phenolic oxygen moiety. The schiff bases  $L^{1}H$  and  $L^{2}H$  show a medium intensity band at around 1327 and 1279 cm<sup>-1</sup> due to phenolic v(C-O) group of is shifted to higher region indicating the coordination through the phenolic oxygen atoms [5]. The IR spectra of the schiff bases  $L^{1}H$  and  $L^{2}H$  exhibit a strong band at 1618 and 1614 cm<sup>-1</sup> due to v(C=N) (azomethine) which has been shifted towards lower region in the spectra of complexes indicating the participation of the azomethine groups in the complex formation [2]. The spectra of the complexes show a broad diffused bands in the region at around 3100-3650 cm<sup>-1</sup>, strong bands at 1535-1538 cm<sup>-1</sup> and week intensity bands at 825-831 cm<sup>-1</sup> due to v(OH),  $\delta$ (OH) and  $\rho_r$ (OH) respectively of the coordinated water molecules [6]. The coordination through nitrogen of azomethine and oxygen of (C-O) group of ligands are further evidenced by the appearance of non-ligand bands in the complexes at around 496-556 cm<sup>-1</sup> and 420-464 cm<sup>-1</sup> are due to M-O and M-N bonds respectively [4], [6].

	IR bands (cm <sup>-1</sup> )							
Schiff base / Complex	v <sub>OH</sub> (phenolic )	ν <sub>OH</sub> (H <sub>2</sub> O)	$\nu_{C=N}$	δ <sub>OH</sub> (H <sub>2</sub> O)	v <sub>C-O</sub> (phenolic)	р <sub>гОН</sub> (H <sub>2</sub> O)	$\nu_{M-O}$	$\nu_{M-N}$
$L^{1}H$	3455	-	1618	-	1327	-	-	-
$L^{2}H$	3447	-	1617	-	1278	-	-	-
$[Co(L^1)(L^2)(H$	-	3150 -	1613,	1534	1387,	825	556,	464,
<sub>2</sub> O) <sub>2</sub> ]		3651	1607		1363		502	417
$[Ni(L^1)(L^2)(H$	_	3200 -	1612,	1533	1384,	826	547,	462,
<sub>2</sub> O) <sub>2</sub> ]		3650	1607	1555	1365	020	502	416
$[Cu(L^1)(L^2)(H$	_	3150 -	1612,	1534	1385,	825	548,	461,
<sub>2</sub> O) <sub>2</sub> ]		3650	1607		1363		496	420
$[Zn(L^1)(L^2)(H$		3100 -	1614,	1538	1393,	831	547,	464,
<sub>2</sub> O) <sub>2</sub> ]	-	3650	1609	1556	1365	031	503	417

			1	2		
Tabla 1.	Chamastanistia	ID hands of th	a liganda I <sup>1</sup>	II and I <sup>4</sup> II and t	hair minad ligand	motol complayer
rable r:	Unaracieristic	TR Dands OF ID	e ngands L	палот, палот	пент пихео прапо	merar complexes.
	enter av ver ibtre	int ound of the	e inganao z		nen minee ngane	metal eompienes.

A4	<b>INTERNATIONAL JOURNAL OF</b>	e-ISSN :	
IJPREMS	PROGRESSIVE RESEARCH IN ENGINEERING	2583-1062	
	MANAGEMENT	<b>.</b>	
	AND SCIENCE (IJPREMS)	Impact	
www.ijprems.com	(Int Peer Reviewed Journal)	Factor : 7.001	
editor@ijprems.com	Vol. 05, Issue 02, February 2025, pp : 1199-1201		

#### Thermogravimetric analysis

The Co(II), Ni(II), Cu(II) and Zn(II) complexes lose their weight in the temperature range ~ 130-272 °C, 125-255 °C, 125-263 °C and 130-276 °C respectively corresponding to two coordinated water molecules with an endothermic peak in DTA curve indicates that the two water molecules are coordinated in the metal complexes [4].

## **4. CONCLUSION**

Magnetic susceptibility, Thermogravimetric analysis and spectral observations suggest the octahedral geometry for the Co(II), Ni(II), Cu(II) and Zn(II) complexes and exhibit coordination number six. The general structure of the complexes is shown in figure 3.



Figure 3: Proposed structure for the complexes.

M = Co(II), Ni(II), Cu(II) and Zn(II).

#### **5. REFERENCES**

- [1] Dash, D.C., Meher F.M., Monhanty P.C. and Nanda J. (1987). Complexes of 3-Benzyllideneimino-2-thiohydantoin and 3-p-Anisylidenemino-2-thiohydantion with Mn(II), Co(II), Ni(II) and Cu(II). Indian Journal Cheemistry, 26(A): 698-701.
- [2] Shriodkar, S. G., Mane, P. S. and Chondhekar, T. K. (2001). Synthesis and fungitoxic studies of Mn(II), Co(II), Ni(II) and Cu(II) with some heterocyclic Schiff base ligands, Indian Journal Chemistry, 40A: 1114-1117.
- [3] Reddy, V., Patil, N. and Angadi, S. D. (2008). Synthesis, Characterization and Antimicrobial Activity of Cu(II), Co(II) and Ni(II) Complexes with O, N, and S Donor Ligands, E-Journal of Chemistry, 5(3):577-583.
- [4] Thakkar, N. V and Bootwala, S. Z. (1995). Synthesis and characterization of binuclear metal complexes derived from some isonitrosoacetophenones and benzidine, Indian Journal Chemistry, 34(A):370-374.
- [5] Makode, J.T. and Aswar, A.S. (2004). Synthesis Characterization, biological and thermal properties of some new Schiff base complexes derived from 2-hydroxy-5-chloroacetophenone and S-methyldithiocarbazate, Indian Journal Cheemistry, 43(A):2120-2125.
- [6] Ravanasiddappa, M., Sureshg, T., Syed, K., Radhavendray, S. C., Basavaraja, C. and Angadi, S., D. (2008). Transition Metal Complexes of 1, 4(2'-Hydroxyphenyl-1-yl) di-iminoazine: Synthesis, Characterization and Antimicrobial Studies, E-Journal of Chemistry, 5(2): 395-403.